

REFERENCES

- CDC. Behavioral Surveillance Workshop. Atlanta: Centers for Disease Control and Prevention; 1995.
- CDC. Guidance. HIV Prevention Community Planning for HIV prevention cooperative agreement recipients. http://www.cdc.gov/nchstp/hiv_aids/pubs/hiv-cp.pdf; 1999.
- Thacker S, Berkelman R. Public health surveillance in the United States. *Epidemiologic Review*. 1988;10:164-190.
- CDC. State-and sex-specific prevalence of selected characteristics--Behavioral Risk Factor Surveillance System, 1994 and 1995. *MMWR*. 1997;46[SS03]:1-29.
- CDC. Trends in sexual risk behaviors among high school students - United States, 1991-1997. *MMWR*. 1998;47:749-752.
- Fowler Jr F. The Redesign of the National Health Interview Survey. *Public Health Rep*. 1996;111:508-11.
- Kelly J, Mosher W, Duffer A, Kinsey S. Plan and operation of the 1995 National Survey of Family Growth. *Vital Health Stat*. 1997;1:1-89.
- Ezzati T, Massey J, Waksberg J, Chu A, Maurer K. Sample design: Third National Health and Nutrition Examination Survey. *Vital Health Stat*. 1992;2:1-35.
- Turner C, Danella R, Rogers R. Sexual behavior in the United States, 1930-1990: trends and methodological problems. *Sex Transm Dis*. 1995;22:173-190.
- Spencer B. On the accuracy of estimates of numbers of intravenous drug users. In: Turner C, Miller H, Moses L, eds. *AIDS Behavior and Intravenous Drug Use*. Washington, DC: National Academy Press; 1989:429-446.
- Michaels S. The prevalence of homosexuality in the United States. In: Cabaj R, Stein T, eds. *Textbook of homosexuality and mental health*. Washington, DC: American Psychiatric Press; 1996:43-63.
- Harrison L. The validity of self-reported drug use in survey research: an overview and critique of research methods. In: Harrison L, Hughs A, eds. *The Validity of Self-Reported Drug Use: Improving the Accuracy of Survey Estimates*; NIDA Research Monograph 167. Rockville, MD: US Department of Health and Human Services; 1997:17-36.
- Bradburn N, Sudman S. *Improving Interview Methods and Questionnaire Design*. San Francisco: Jossey-Bass; 1979.

CDC. Update: trends in AIDS incidence, deaths, and prevalence - United States, 1996. MMWR. 1997;46:165-173.

Anderson J. CDC data systems collecting behavioral data on HIV counseling and testing. Public Health Rep. 1996;111(S-1):129-132.

CDC. HIV/AIDS Surveillance Report. Atlanta, GA: Centers for Disease Control and Prevention; 1999.

Buehler J, Diaz T, Hersh B, Chu S. The Supplement to HIV/AIDS Surveillance Project: an approach for monitoring HIV risk behaviors. Public Health Rep. 1996;111(S-1):133-137.

Rietmeijer C, Davidson A, Cohn D. Substance use and high-risk sexual behaviors among persons with HIV infection accessing medical services in a public health care system. . 1999 National HIV Prevention Conference. Atlanta, GA; 1999:Abstract #226.

Campsmith M, Nakashima A, Jones J, the SHAS Project Group. Does crack use after HIV diagnosis predict high risk sexual behaviors? Differences between men and women. 1999 National HIV Prevention Conference. Atlanta, GA; 1999:Abstract #159.

Nakashima A, Burgess D, Campsmith M, et al. Representativeness of HIV/AIDS cases interviewed in the Supplement to HIV/AIDS Surveillance (SHAS) Project. 1999 National HIV Prevention Conference. Atlanta, GA; 1999:Abstract #160.

CDC. HIV testing among populations at high risk for HIV infection - nine states, November 1995 - December 1996. MMWR. 1998;47:87-100.

Wortley P, Lehman J, Fleming P. Name-based reporting of HIV infection. Am J Public Health. 1999;89:1271-1272.

McKellar D, Valleroy L, Karon J, Lemp G, Janssen R. The young men's survey: methods for estimating HIV seroprevalence and risk factors among young men who have sex with men. Public Health Rep. 1996;111(S-1):138-144.

Valleroy L, McKellar D, Rosen D, Secura G. Prevalence and predictors of unprotected receptive anal intercourse for 15- to 22-year-old men who have sex with men in seven urban areas, USA. XII International Conference on AIDS. Geneva, Switzerland; 1998:Abstract #23139.

CDC Advisory Committee on the Prevention of HIV Infection. External review of CDC's HIV prevention activities. Atlanta: Centers for Disease Control and Prevention; 1995.

Anderson J, Rietmeijer C, Wilson R, Barker P. Asking about condom use: is there a standard approach that should be adopted across surveys? Proceedings of the American Statistical Association. 1998.

Fishbein M. Toward an understanding of the relationships between behavioral and biological outcome measures. 13th Meeting of the International Society for Sexually Transmitted Diseases Research. Denver, CO, USA; 1999.

Dillon B, Hecht F, Swanson M, Goupil-Sormany I, Grant R. Primary HIV infections associated with oral transmission. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco; 2000:Abstract #473.

Rietmeijer C, Fichtner R, and the Behavioral Surveillance Workgroup. Use of condoms at last sexual intercourse: an example of advantages and pitfalls of standardizing questions in surveys for HIV and STD risk behavior. XII International Conference on AIDS. Geneva, Switzerland; 1998: Abstract #43427.

Leonard L, Ross M. The last sexual encounter: the contextualization of sexual risk behavior. Int J STD AIDS. 1997;8:643-5.